

## Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004-2005. For more information, please visit [www.landfire.gov](http://www.landfire.gov). Please direct questions to [helpdesk@landfire.gov](mailto:helpdesk@landfire.gov).

### Potential Natural Vegetation Group (PNVG):

R2ASMC1w

Aspen with Conifer--Low to Mid-Elevations

### General Information

**Contributors** (additional contributors may be listed under "Model Evolution and Comments")

#### Modelers

Linda Chappell      lchappell@fs.fed.us  
  
Bob Campbell      rbcampbell@fs.fed.us  
Cheri Howell      chowell02@fs.fed.us

#### Reviewers

Krista Gollnick-      Krista\_Waid@blm.gov  
Wade/Sarah Heidi  
Charles E. Kay      ckay@hass.usu.edu  
Wayne D. Shepperd      wshepperd@fs.fed.us

#### Vegetation Type

Forested

#### Dominant Species\*

POTR5  
PIPO  
PICO  
PSME

#### General Model Sources

- Literature  
 Local Data  
 Expert Estimate

#### LANDFIRE Mapping Zones

12	17
13	18
16	

#### Rapid Assessment Model Zones

- |   |  |
|---|--|
| <input type="checkbox"/> California             | <input type="checkbox"/> Pacific Northwest |
| <input checked="" type="checkbox"/> Great Basin | <input type="checkbox"/> South Central     |
| <input type="checkbox"/> Great Lakes            | <input type="checkbox"/> Southeast         |
| <input type="checkbox"/> Northeast              | <input type="checkbox"/> S. Appalachians   |
| <input type="checkbox"/> Northern Plains        | <input type="checkbox"/> Southwest         |
| <input type="checkbox"/> N-Cent.Rockies         |  |

### Geographic Range

Great Basin, California, northern Rockies, Alaska, Pacific Northwest, and north central regions.

### Biophysical Site Description

This type typically occurs on flat to steep terrain (<80%) on all aspects. Elevation ranges from 5000' to 9000'. Higher latitude or northern aspects tend toward the lower elevation range while lower latitudes and southern aspects tend toward the higher elevation range. Soils are highly variable, but generally cool. This type occurs above the pinyon/juniper and/or sagebrush but below the spruce-fir cover.

### Vegetation Description

Without regular fire and with high levels of herbivory, conifers may replace the aspen community. The presence of even a single aspen tree in a stand provides strong evidence that the area historically supported an aspen cover type. Areas with as few as five aspen trees per acre may return to an aspen community following disturbance.

As a species, aspen is adapted to a much broader range of environments than most plants found associated with it. Aspen exists in single-storied or multi-storied stands. Conifer species are common and often include Douglas-fir (*Pseudotsuga menziesii*), white fir (*Abies concolor*), ponderosa pine (*Pinus ponderosa*) and/or lodgepole pine (*Pinus contorta*). Historically ponderosa pine was the fire adapted species that occurred in open savannahs with old ponderosa pine on the ridges or rocky outcrops that provided some protection from periodic fire. Aspen could function as a tall shrub rather than an overstory tree because of fire's frequent return.

### Disturbance Description

This is a strongly fire adapted community, with FRIs varying greatly with the encroachment of conifers.

\*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Some sites are prone to snowslides, mudslides and rotational slumping. Flooding may also operate in these systems. Before conifer encroachment in developing stands (<40 yrs), we adopted the FRI of stable aspen (R2ASPN), i.e., no fire in early development and only replacement fire every 75-yr in young stand between 10-40 yrs old. Similarly, older stands dominated by conifers would experience replacement fire every 75 yrs. For stands between 40-125 yrs with encroaching conifers, replacement, mixed severity, and surface fires were more frequent. According to Baker (1925), who most closely studied the historic condition, the FRI for replacement fire was 20-40 yrs (min-max). The FRI for mixed severity fire and surface fire was 10-20 yrs (min-max). Mixed severity fire was found in closed stands between 40-125 yrs, whereas surface fire was found in open stand >40 yrs, which were less common, based on frequent fire scars left on aspen. Indian burning was the primary sources of fire, especially surface fire. Studies by Bartos and Campbell (1998) support these findings. It is important to understand that aspen is considered a fire-proof vegetation type that does not burn during the normal lightning season, yet evidence of frequent fire scars and historical studies show that native burning was the only source of fire that occurred mostly during the spring and fall.

### Adjacency or Identification Concerns

This includes low elevation lodgepole, not the subalpine-fir mix.

If conifers are not present, the stable aspen model should be considered. If subalpine fir or spruce are present, the aspen w/mixed conifers for high elevation model should be considered.

### Scale Description

Sources of Scale Data  Literature  Local Data  Expert Estimate

This type occurs in a landscape mosaic from moderate to large sized patches.

### Issues/Problems

There is uncertainty about the role of mixed severity fire. We assumed that native burning in aspen stands invaded by young conifers resulted in mixed severity fire, whereas the same source of fire would cause low severity fire (surface fire) in same age stands that were more open. Experts and modelers expressed different views about the frequency of all fires, citing FRIs longer than those noted by Baker (1925), who actually studied the historic condition. The FRIs used here were a compromise: 1) the longer FRIs were used for the earlier and oldest development states and 2) the maximum FRI of Baker (1925) was used for stands between 40 and 125 yrs that were being encroached by lower elevation conifers.

### Model Evolution and Comments

This type is more highly threatened by conifer replacement than stable aspen. As this type has a fairly short fire return interval compared to other aspen types, it should be noted that aspen can act as a tall shrub. Bradley, et. Al. (1992) state that Loope & Gruell estimated a fire frequency of 25 to 100 years for a Douglas-fir forest with seral aspen in Grand Teton National Park (p39). They later state that fire frequencies of 100 to 300 years appear to be appropriate for maintaining most seral aspen stands. In the Fontenelle Creek, Wyoming drainage, the mean fire-free interval was estimated to be 40 years. Fires in this area burned in a mosaic pattern of severities, from stand-replacement to low fires that scarred but did not kill the relatively thin-barked lodgepole pine on the site (p46).

## Succession Classes

*Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook ([www.frcc.gov](http://www.frcc.gov)).*

**Class A 15%**

Early1 PostRep

**Description**

Grass/forb and aspen suckers <6'tall. Generally, this is expected to occur 1-3 years post-disturbance. Replacement fire was absent. Succession to B after 10 years.

**Indicator Species\* and Canopy Position**

POTR5

**Upper Layer Lifeform**

- Herbaceous  
 Shrub  
 Tree

**Fuel Model** no data**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	0 %	99 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class B 40%**

Mid1 Closed

**Description**

Aspen saplings over 6' tall dominate. Canopy cover is highly variable. Conifers can invade. The stand is composed of 80% aspen, up to 10 % conifers. The FRI of 75 yrs was used for replacement fire. Succession to C after 30 years.

**Indicator Species\* and Canopy Position**

POTR5

PIPO

PICO

**Upper Layer Lifeform**

- Herbaceous  
 Shrub  
 Tree

**Fuel Model** no data**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	20 %	99 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class C 15%**

Mid2 Closed

**Description**

Aspen 5" to 16". Mixed aspen overstory and conifer understory dominance. Less than 25 % conifer. Native burning causes both replacement fire (using Baker's max FRI of 40 yrs) and mixed severity fire (using Baker's max FRI of 20 yrs) that opens this stand by thinning conifers and aspen (disturbance to D). In the absence of fire it will naturally succeed to a closed conifer stand (E) after 85 years.

**Indicator Species\* and Canopy Position**

POTR5

PIPO

PICO

PSME

**Upper Layer Lifeform**

- Herbaceous  
 Shrub  
 Tree

**Fuel Model** no data**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	20 %	99 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class D 25 %**

Late1 Open

**Description**

Aspen dominate with conifer understory up to co-dominance: 80% aspen overstory. Conifers (e.g., ponderosa pine) are assumed more resistant to fire than aspen and will likely cause the progressive suppression of aspen. Surface fire keeps this stand open, kills young conifers, and maintains aspen: every 20 yrs (max FRI from Baker). Replacement fire is every 40 years, the maximum from Baker. In the absence of any fire for 2-3 FRIs (84 yrs), the stand will become closed with conifers (E).

**Indicator Species\* and Canopy Position**

POTR5  
ABCO

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	0 %	39 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class E 5 %**

Late1 Closed

**Description**

Conifers dominate at 125+ years. Aspen over 16", mixed conifer mixed sizes, main overstory is conifers. Greater than 50% conifer in the overstory. FRI for replacement fire is every 75 years.

**Indicator Species\* and Canopy Position**

PIPO  
PICO  
PSME  
POTR5

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	40 %	80 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Disturbances**

**Non-Fire Disturbances Modeled**

- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other:

**Fire Regime Group: 2**

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

**Historical Fire Size (acres)**

Avg:  
Min:  
Max:

**Fire Intervals (FI):**

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

\*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

<b>Sources of Fire Regime Data</b>	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
	<i>Replacement</i>	61		0.01639	53
<input checked="" type="checkbox"/> Literature	<i>Mixed</i>	137		0.0073	24
<input checked="" type="checkbox"/> Local Data	<i>Surface</i>	143		0.00699	23
<input checked="" type="checkbox"/> Expert Estimate	<i>All Fires</i>	33		0.03069	

## **References**

Baker, F. S., 1925. Aspen in the Central Rocky Mountain Region. USDA Department Bulletin 1291 pp. 1-47.

Bartos, D. L. 2001. Landscape Dynamics of Aspen and Conifer Forests. Pages 5-14 in: Shepperd, W. D.; Binkley, D.; Bartos, D. L.; Stohlgren, T. J.; and Eskew, L. G., compilers. 2001. Sustaining aspen in western landscapes: symposium proceedings; 13-15 June 2000; Grand Junction, CO. Proceedings RMRS-P-18. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 460 p.

Bartos, D. L. and R. B. Campbell, Jr. 1998. Decline of Quaking Aspen in the Interior West – Examples from Utah. *Rangelands*, 20(1):17-24.

Bradley, A. E., Noste, N. V., and W. C. Fischer. 1992. Fire Ecology of Forests and Woodlands in Utah. GTR-INT-287. Ogden, UT. U.S. Department of Agriculture, Forest Service, Intermountain Research Station. 128 p.

Bradley, Anne E., W. C. Fischer, and N. V. Noste. 1992. Fire Ecology of the Forest Habitat Types of Eastern Idaho and Western Wypoming. GTR- INT-290. Ogden, UT. U.S. Department of Agriculture, Forest Service, Intermountain Research Station. 92.

Brown, J.K. and D.G. Simmerman. 1986. Appraisal of fuels and flammability in western aspen: a prescribed fire guide. General technical report INT-205. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.

Brown, J. K., K. Smith, J. Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Campbell, R. B. and , D. L. Bartos. 2001. Objectives for Sustaining Biodiversity. In: Shepperd, W. D.; Binkley, D.; Bartos, D. L.; Stohlgren, T. J.; and Eskew, L. G., compilers. 2001. Sustaining aspen in western landscapes: symposium proceedings; 13-15 June 2000; Grand Junction, CO. Proceedings RMRS-P-18. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 460 p.

Debyle, N.V., C.D. Bevins, and W.C. Fisher. 1987. Wildfire occurrence in aspen in the interior western United States. *Western Journal of Applied Forestry*. 2:73-76.

Kay, C. E. 1997. Is aspen doomed? *Journal of Forestry* 95: 4-11.

Kay, C. E. 2001. Evaluation of burned aspen communities in Jackson Hole, Wyoming. Proceedings RMRS-P-18. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 8 p.

Kay, C.E. 2001. Long-term aspen exclosures in the Yellowstone ecosystem. Proceedings RMRS-P-18.. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 15 p.

Kay, C.E. 2001. Native burning in western North America: Implications for hardwood forest management. General Technical Report NE-274. U.S. Department of Agriculture, Forest Service, Northeast Research Station. 8 p.

Mueggler, W. F. 1988. Aspen Community Types of the Intermountain Region. USDA Forest Service, General Technical Report INT-250. 135 p.

Mueggler, W. F. 1989. Age Distribution and Reproduction of Intermountain Aspen Stands. Western Journal of Applied Forestry, 4(2):41-45.

Romme, WH, L. Floyd-Hanna, D. D. Hanna ,and E. Bartlett. 2001. Aspen's ecological role in the west. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, RMRS Proceedings-P-18. Pages 243-259.

Shepperd, W.D. and E.W. Smith. 1993. The role of near-surface lateral roots in the life cycle of aspen in the central Rocky Mountains. Forest Ecology and Management 61: 157-160.

Shepperd, W. D. 2001. Manipulations to Regenerate Aspen Ecosystems. Pages 355-365 in: Shepperd, Wayne D.; Binkley, Dan; Bartos, Dale L.; Stohlgren, Thomas J.; and Eskew, Lane G., compilers. 2001. Sustaining aspen in western landscapes: symposium proceedings; 13-15 June 2000; Grand Junction, CO. Proceedings RMRS-P-18. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 460 p.

Shepperd, W. D., D. L. Bartos, and A. M. Stepen. 2001. Above- and below-ground effects of aspen clonal regeneration and succession to conifers. Canadian Journal of Forest Resources; 31: 739-745.

USDA Forest Service. 2000. Properly Functioning Condition: Rapid Assessment Process (January 7, 2000 version). Intermountain Region, Ogden, UT. Unnumbered.

Welsh, S. L, N. D. Atwood, S. I. Goodrich, and L. C. Higgins. 2003. A Utah Flora, Third edition, revised. Print Services, Brigham Young University, Provo, UT. 912 p.